



This form should be used for all taxonomic proposals. Please complete all those modules that are applicable (and then delete the unwanted sections). For guidance, see the notes written in blue and the separate document "Help with completing a taxonomic proposal"

Please try to keep related proposals within a single document; you can copy the modules to create more than one genus within a new family, for example.

MODULE 1: **TITLE, AUTHORS, etc**

Code assigned:	<i>2013.013a-dV</i>	(to be completed by ICTV officers)			
Short title: Create a new species, <i>Pasivirus A</i> , in a new genus, <i>Pasivirus</i> , within the family <i>Picornaviridae</i> (order <i>Picornavirales</i>) (e.g. 6 new species in the genus <i>Zetavirus</i>)					
Modules attached (modules 1 and 9 are required)	1 <input checked="" type="checkbox"/> 6 <input type="checkbox"/>	2 <input checked="" type="checkbox"/> 7 <input type="checkbox"/>	3 <input checked="" type="checkbox"/> 8 <input type="checkbox"/>	4 <input type="checkbox"/> 9 <input checked="" type="checkbox"/>	5 <input type="checkbox"/>

Author(s) with e-mail address(es) of the proposer:

Nick J. Knowles (nick.knowles@pirbright.ac.uk) on behalf of the *Picornaviridae* Study Group

List the ICTV study group(s) that have seen this proposal:

A list of study groups and contacts is provided at <http://www.ictvonline.org/subcommittees.asp> . If in doubt, contact the appropriate subcommittee chair (fungal, invertebrate, plant, prokaryote or vertebrate viruses)

Picornaviridae Study Group

ICTV-EC or Study Group comments and response of the proposer:

Date first submitted to ICTV:

25/06/2013

Date of this revision (if different to above):

MODULE 2: **NEW SPECIES**

creating and naming one or more new species.

If more than one, they should be a group of related species belonging to the same genus. All new species must be placed in a higher taxon. This is usually a genus although it is also permissible for species to be “unassigned” within a subfamily or family. Wherever possible, provide sequence accession number(s) for one isolate of each new species proposed.

Code	2013.013aV	(assigned by ICTV officers)
To create one new species within:		
Genus:	<i>Pasivirus</i> (new)	Fill in all that apply. • If the higher taxon has yet to be created (in a later module, below) write “ (new) ” after its proposed name. • If no genus is specified, enter “ unassigned ” in the genus box.
Subfamily:	n/a	
Family:	<i>Picornaviridae</i>	
Order:	<i>Picornavirales</i>	
And name the new species:		GenBank sequence accession number(s) of reference isolate:
<i>Pasivirus A</i>		JQ316470, JX491648

Reasons to justify the creation and assignment of the new species:

- Explain how the proposed species differ(s) from all existing species.
 - If species demarcation criteria (see module 3) have previously been defined for the genus, **explain how the new species meet these criteria.**
 - If criteria for demarcating species need to be defined (because there will now be more than one species in the genus), please state the proposed criteria.
- Further material in support of this proposal may be presented in the Appendix, Module 9

Virus discovery

Sauvage et al. (2012) recently described the molecular detection of a new picornavirus in healthy pigs in France. A second swine pasivirus genome sequence (from pig faeces) was submitted to GenBank on 15-AUG-2012 (Duan and Yu, unpub.).

Growth in cell cultures

The virus has not been cultivated in cell cultures.

Untranslated regions

Only a partial sequence (378 nt) of the 5' UTR is available and the internal ribosome entry site (IRES) type is unknown. The 3' UTR is 116 nt long. Neither region has any significant identity with other picornaviruses.

Genome organization/proteins

VPg+5'UTR[1AB-1C-1D-2A1^{npgp}/2A2-2B-2C/3A-3B^{VPg}-3C^{pro}-3D^{pol}]3'UTR-poly(A)

[], defines the long ORF encoding the polyprotein.

/, Indicates primary polyprotein cleavages.

-, indicates secondary cleavages mainly performed by the 3C^{pro} polypeptide.

VP0 (1AB) is predicted to not be cleaved to VP4/VP2 and does not have a myristoylation

signal (GxxxT/S) at its amino-terminus. 2A1 is either very short or is part of VP1. 2A2 is 112 aa in length and has no similarity to other picornavirus proteins.

Genetic relationships

The two pasivirus genomes share 80% nt identity. Their VP1 nt and aa sequences differ by about 24.5% and 16.5%, respectively; therefore they may represent two distinct (geno)types. Pasiviruses are most closely related to the members of the *Parechovirus* genus in the P1 (*Ljungan virus*; 35.4%), P2 (*Human parechovirus*, 34.8%) and P3 (*Ljungan virus*; 32.5%) polypeptides.

MODULE 3: **NEW GENUS**

creating a new genus

Ideally, a genus should be placed within a higher taxon.

Code	2013.013bV	(assigned by ICTV officers)
To create a new genus within:		
Subfamily:	n/a	Fill in all that apply. • If the higher taxon has yet to be created (in a later module, below) write “ (new) ” after its proposed name. • If no family is specified, enter “ unassigned ” in the family box
Family:	Picornaviridae	
Order:	Picornavirales	

naming a new genus

Code	2013.013cV	(assigned by ICTV officers)
To name the new genus: <i>Pasivirus</i>		

Assigning the type species and other species to a new genus

Code	2013.013dV	(assigned by ICTV officers)
To designate the following as the type species of the new genus		
<i>Pasivirus A</i>	Every genus must have a type species. This should be a well characterized species although not necessarily the first to be discovered	
The new genus will also contain any other new species created and assigned to it (Module 2) and any that are being moved from elsewhere (Module 7b). Please enter here the TOTAL number of species (including the type species) that the genus will contain:		
1		

Reasons to justify the creation of a new genus:

Additional material in support of this proposal may be presented in the Appendix, Module 9

Pasiviruses have a similar genome organization to the parechoviruses, but differ in that the parechovirus 2A contains an H-box/NC motif. Pasiviruses are also most closely related to the members of the *Parechovirus* genus in the P1 (*Ljungan virus*; 35.4%), P2 (*Human parechovirus*, 34.8%) and P3 (*Ljungan virus*; 32.5%) polypeptides. However, the *Picornaviridae* Study Group (PSG) guidelines state that members of different genera share less than 40%, 40% and 50% amino acid difference in P1, P2 and P3, respectively. We therefore suggest that the proposed species *Pasivirus A* is placed in a new genus named *Pasivirus*.

Origin of the new genus name:

Pasivirus, from **Parecho** sister-clade virus

Reasons to justify the choice of type species:

The genus is proposed to contain only a single species.

Species demarcation criteria in the new genus:

If there will be more than one species in the new genus, list the criteria being used for species demarcation and explain how the proposed members meet these criteria.

None, since there is only a single species.

MODULE 9: **APPENDIX**: supporting material

additional material in support of this proposal

References:

Sauvage, V., Ar Gouilh, M., Cheval, J., Muth, E., Pariente, K., Burguiere, A., Caro, V., Manuguerra, J.C. and Eloit, M. (2012). A member of a new *Picornaviridae* genus is shed in pig feces. *J. Virol.* 86: 10036-10046. Epub 2012 Jul 11.

Yu, J.M., Li, J.S., Ao, Y.Y. and Duan, Z.J. (2013). Detection of novel viruses in porcine fecal samples from China. *Virol J.* 10(1): 39 [Epub 2013 Jan 30].

Annex:

Include as much information as necessary to support the proposal, including diagrams comparing the old and new taxonomic orders. The use of Figures and Tables is strongly recommended but direct pasting of content from publications will require permission from the copyright holder together with appropriate acknowledgement as this proposal will be placed on a public web site. For phylogenetic analysis, try to provide a tree where branch length is related to genetic distance.




Figure 1. Maximum likelihood tree showing the relationship between picornaviruses in the P1 capsid. Sequences were aligned using MUSCLE and the tree constructed using MEGA 5.2.




Figure 2. Maximum likelihood tree showing the relationship between picornaviruses in the 3D polymerase. Sequences were aligned using MUSCLE and the tree constructed using MEGA 5.2.